



## Application of digital image treatment to the characterization and differentiation of deep-sea ichnofacies

Javier DORADOR\* & Francisco J. RODRÍGUEZ-TOVAR

Departamento de Estratigrafía y Paleontología, Universidad de Granada, 18002 Granada, Spain; javidr@ugr.es; fjrtovar@ugr.es

\* Corresponding author

Dorador, J. & Rodríguez-Tovar, F.J. 2015. Application of digital image treatment to the characterization and differentiation of deep-sea ichnofacies. [Aplicación del tratamiento de imagen a la caracterización y diferenciación de icnofacies marinas profundas]. *Spanish Journal of Palaeontology*, 30 (2), 265-274.

Manuscript received 20 December 2014  
Manuscript accepted 01 July 2015

© Sociedad Española de Paleontología ISSN 2255-0550

### ABSTRACT

The ichnofacies model proposed by Seilacher in the 1960's stands as one of the main tools used in the ichnological research. Based on the integration of numerous ichnological and sedimentological observations, it entails a precise characterization of the trace fossil assemblage. Differentiation of trace fossils may be relatively easy in outcrops, but cores are a different matter, particularly modern marine cores. Difficulties are even more accentuated in deep-sea pelagic and hemipelagic, non-turbiditic, fine-grained deposits. The application of digital image treatment can facilitate trace fossil identification and improve deep-sea ichnofacies characterization on modern cores. The method proposed here was applied to cores from the IODP Expedition 339 (Site U1385). Eight ichnotaxa —*Chondrites*, *Palaeophycus*, *Phycosiphon*, *Planolites*, *Taenidium*, *Thalassinoides*, *Thalassinoides*-like structures and *Zoophycos*— as well as some horizontal grouped circular structures interpreted as trails (?*Nereites*) were recognized. The archetypical *Zoophycos* ichnofacies is identified, but characteristic elements belonging to the *Nereites* ichnofacies and to the distal expression of *Cruziana* ichnofacies are locally recognized. Thus, the applied methodology allows differentiating between different

### RESUMEN

El modelo de icnofacies, propuesto por Seilacher en la década de los 60, es una de las principales herramientas aplicadas en icnología. Basado en la integración de numerosas observaciones icnológicas y sedimentológicas, una precisa caracterización de la asociación de icnotaxones es esencial en el análisis de icnofacies. La diferenciación de icnotaxones puede ser relativamente fácil en afloramiento, pero es comparativamente difícil en sondeos, especialmente si corresponden a materiales marinos modernos. La dificultad se acentúa cuando se trata de sedimentos pelágicos, hemipelágicos, no turbidíticos, de grano fino. Se propone la aplicación del tratamiento de imágenes para facilitar la identificación de icnotaxones y mejorar la caracterización de icnofacies marinas profundas en sondeos modernos. El método ha sido aplicado a sondeos de la Expedición 339 del IODP (Site U1385). Ocho icnotaxones, *Chondrites*, *Palaeophycus*, *Phycosiphon*, *Planolites*, *Taenidium*, *Thalassinoides*, estructuras similares a *Thalassinoides* y *Zoophycos*, y algunas formas circulares, densamente agrupadas, interpretadas como rastros horizontales (?*Nereites*), han sido reconocidas. Se ha identificado claramente la icnofacies de *Zoophycos*, aunque se reconocen elementos propios de las icnofacies de

expressions of the *Zoophycos* ichnofacies and to approach subtle changes in the involved environmental parameters affecting tracemakers.

**Keywords:** Trace fossils, ichnofacies, deep-sea, IODP Expedition 339, Site U1385, digital image treatment.

*Nereites* y de la expresión distal de icnofacies de *Cruziana*. La metodología aplicada facilita el reconocimiento de distintas expresiones de la icnofacies de *Zoophycos* y permite interpretar sutiles cambios en los parámetros ambientales que afectan a los bioturbadores.

**Palabras clave:** Icnofósiles, icnofacies, medios marinos profundos, Expedición 339 IODP, Site U1385, tratamiento digital de imágenes.

## 1. INTRODUCTION

The ichnofacies model, proposed by Seilacher during the 1960's (Seilacher, 1964, 1967), led to unprecedented development of ichnological analysis. This model is based on the identification of certain features shared by ichnotaxa during a longtime span that can be related to particular environmental conditions (Buatois & Mángano, 2011). Initially just six ichnofacies were proposed, but over the past five decades the model has been modified and expanded to embrace many more recurrent, archetypal (Seilacherian) ichnofacies that play an important role in basin analysis (see recent reviews in Buatois & Mángano, 2011; MacEachern *et al.*, 2010, 2012). To date, 15 archetypal marine and continental ichnofacies have been defined based on invertebrate trace fossil assemblages (Buatois & Mángano, 2011). Earlier papers focused on application of ichnofacies as palaeobathymeters (e.g., Seilacher, 1967; Frey & Pemberton, 1985; Bottjer *et al.*, 1987). However, palaeobathymetry is not the only controlling factor —ichnofacies depend on the interplay of many parameters, such as substrate, energy, food supply, oxygenation level and taphonomic features that control the distribution of tracemakers and determine the preservation of biogenic structures (e.g., Bromley, 1990, 1996; Wetzel, 1991; Buatois & Mángano, 1995; MacEachern *et al.*, 2007). Accordingly, detailed ichnofacies characterization has come to be an essential tool for palaeoenvironmental research, the interpretation of depositional settings, and basin analysis.

Ichnofacies characterization is not a mere listing of ichnotaxa. Rather, it is based on the integration of a complete data set that integrates ichnologic, sedimentologic, stratigraphic and taphonomic information. Still, precise ichnotaxonomic differentiation is fundamental for ichnofacies analysis, providing important palaeoecologic information (Wetzel, 1991). Such differentiation is relatively easy in outcrops, but complicated when working on core materials, owing to narrow views of, limited size and almost exclusive two-dimensional expression of trace fossils (e.g., Bromley, 1996; Wetzel, 2010; Dorador *et al.*, 2014a). The difficulty is even greater when working with modern marine cores, especially those of deep-sea,

pelagic-hemipelagic sediments in which trace fossils have diffuse boundaries and their infilling materials are visually very similar to the host sediment.

Several techniques have been developed to improve ichnotaxonomic differentiation in modern marine cores. Some of them, for instance X-ray scanning (e.g., Howard, 1968; Löwemark & Werner, 2001) and computed tomography (Joschko *et al.*, 1991; Rosenberg *et al.*, 2007, among others), require specialized methodology and instrumentation, making them comparatively complex, expensive and infrequently employed. Magwood & Ekdale (1994) used image treatment for ichnotaxa differentiation and ichnofabric visualization by means of matrix operations. However, the complexity of its application is an impediment for the widespread use of this method. Recently, an inexpensive high-resolution image treatment method that is easy to use was developed for the ichnological study of modern deep-sea marine cores. The advantages of this novel technique are: 1) differentiation between biodeformational structures and trace fossils (Dorador *et al.*, 2014a), 2) quantification of the percentage of bioturbation in an objective and semi-automatic way (Dorador *et al.*, 2014b), 3) knowledge of overlapping between trace fossils, tiering structure and penetration depth of some ichnotaxa (Dorador & Rodríguez-Tovar, 2014), and 4) enhancement of ichnofabric characterization (Rodríguez-Tovar & Dorador, 2014, 2015). This paper describes the usefulness of this high-resolution image technique for the ichnotaxonomic characterization of trace fossil assemblages pertaining to deep-sea ichnofacies in cores of modern marine deposits.

### 1.1. Deep-marine ichnofacies

Deep-marine sediments are mainly characterized by the *Zoophycos* and *Nereites* ichnofacies (Uchman & Wetzel, 2011; Wetzel & Uchman, 2012; Fig. 1).

The *Zoophycos* ichnofacies traditionally has been located between the *Cruziana* and *Nereites* ichnofacies, below storm-wave base in outer shelf to slope deposits (Seilacher, 1967; Uchman & Wetzel, 2011). Hence, it is considered as intermediate between ichnofacies typical

of inner shelf or offshore deposits (i.e., the *Cruziana* ichnofacies) and those of abyssal deposits (i.e., the *Nereites* ichnofacies) (Hubbard *et al.*, 2012). The *Zoophycos* ichnofacies is usually associated with fine-grained, pelagic and hemipelagic, non-turbiditic sediments, and with very low sedimentation rates. It is commonly linked to low-energy subtidal settings, periods of dysaerobic or anoxic conditions, and sediments with abundant organic matter. In this context, diversity is usually low and abundance is high, and the trace fossil assemblage is characterized by a prevalence of relatively simple to complex feeding structures with spreite, subordinate grazing traces, and the dominance of deep-tier forms of deposit feeders and farmers. One typical ichnogenus is *Zoophycos*, but *Chondrites* and *Phycosiphon* are also characteristic. *Cosmorhapha*, *Helminthopsis*, *Planolites*, *Scolicia*, *Spirophyton* and *Thalassinoides* are also present, as well as certain ichnospecies of *Nereites* (Buatois & Mángano, 2011; MacEachern *et al.*, 2012; Uchman & Wetzel, 2012; Table 1).


**Table 1.** Common trace fossils registered in the distal expression of *Cruziana* ichnofacies, *Zoophycos* ichnofacies and *Nereites* ichnofacies. Note: Grey tone for more (dark) or less (light) abundant ichnotaxa.

Ichnotaxa	Distal expression of <i>Cruziana</i> ichnofacies	<i>Zoophycos</i> ichnofacies	<i>Nereites</i> ichnofacies
<i>Asterosoma</i>	—	—	—
<i>Chondrites</i>	—	—	—
<i>Cosmorhapha</i>	—	—	—
<i>Cylindrichnus</i>	—	—	—
<i>Graphoglyptids</i>	—	—	—
<i>Helminthopsis</i>	—	—	—
<i>Nereites</i>	—	—	—
<i>Ophiomorpha</i>	—	—	—
<i>Phycosiphon</i>	—	—	—
<i>Planolites</i>	—	—	—
<i>Rosselia</i>	—	—	—
<i>Scolicia</i>	—	—	—
<i>Spirophyton</i>	—	—	—
<i>Thalassinoides</i>	—	—	—
<i>Zoophycos</i>	—	—	—

The *Nereites* ichnofacies is common in basin-floor depositional environments characterized by very slow pelagic and hemipelagic background sedimentation interrupted by periodic turbidite deposition. Ichnodiversity is very high and abundance is high. Trace fossil assemblages are characterized by dominance of complex graphoglyptids

reflecting farming or trapping (in response to the slow accumulation of food), and the presence of grazing trails and feeding traces of detritus and deposit feeders; shallow-tier trace fossils are generally dominant (Buatois & Mángano, 2011; MacEachern *et al.*, 2012; Uchman & Wetzel, 2012). Taphonomy plays an important role in this ichnofacies, particularly in the preservation of delicate graphoglyptid structures (Uchman & Wetzel, 2012). This ichnofacies can be divided into three ichnosubfacies (Seilacher, 1974; Uchman, 2009) to distinguish the assemblages related to different parts of a turbidite (channel, lobes, etc.): 1) The *Ophiomorpha rudis* ichnosubfacies is related to thick-bedded sandstones in turbidite successions in channels or proximal lobes, and mainly consists of *Ophiomorpha*, *Scolicia*, and occasionally *Nereites* and *Chondrites* (Uchman, 2009; Buatois & Mángano, 2011); 2) the *Paleodictyon* ichnosubfacies is characterized by sandier, medium- to thin- bedded flysch deposits, featuring the presence of graphoglyptids; and 3) the *Nereites* ichnosubfacies represents the most distal part of turbidites (muddy distal flysch sediments) and is characterized by the presence of backfilled trace fossils such as *Nereites*, *Phycosiphon* or *Zoophycos* (Seilacher, 1974; Heard & Pickering, 2008; Knaust, 2009; Uchman, 2009; Buatois & Mángano, 2011; Uchman & Wetzel, 2012; Table 1). The succession of these three ichnosubfacies can be related to bathymetric trends from shallow to deeper parts, and with non-bathymetric trends from channel axis, levee to overbank or inter-channel areas (Uchman & Wetzel, 2012).

Additionally, the distal expression of the *Cruziana* ichnofacies can be identified together with the *Zoophycos* and *Nereites* ichnofacies in deep-sea settings. The archetypal *Cruziana* ichnofacies occurs from just slightly above fair-weather base to the storm-wave base, from the lower shoreface to the lower offshore, under moderate- to low-energy conditions. The wide range of zones, implying variations in palaeoenvironmental conditions and, hence, ichnological assemblages, allows for subdivisions of this archetypal *Cruziana* ichnofacies (MacEachern *et al.*, 1999, 2007). The distal expression of the *Cruziana* ichnofacies is transitional between the archetypal *Cruziana* ichnofacies and the archetypal *Zoophycos* ichnofacies in a basinward direction, usually from lower offshore toward inner shelf, and is associated with muddy siltstones and silty mudstones (MacEachern *et al.*, 1999, 2007; Buatois & Mángano, 2011). This distal expression of the *Cruziana* ichnofacies is dominated by feeding and grazing structures of deposit feeders, with suspension-feeding structures generally absent. There is an overall decrease in *Rosselia*, *Cylindrichnus*, *Asterosoma* and *Ophiomorpha*, with an increase in *Helminthopsis*, *Phycosiphon*, *Planolites*, *Thalassinoides* and *Zoophycos*, compared to the archetypal expression of the *Cruziana* ichnofacies (MacEachern *et al.*, 1999, 2007; Buatois & Mángano, 2011; Table 1).



**Figure 1.** Schematic ichnofacies diagram showing the distribution of the most common ichnogenera along a proximal-distal trend, from sublittoral to abyssal zone. Modified from Buatois & Mángano (2011) and Pemberton *et al.* (2012). *As*, *Asterosoma*; *Ch*, *Chondrites*; *Co*, *Cosmorhaphis*; *Cy*, *Cylindrichnus*; *He*, *Helminthopsis*; *Ne*, *Nereites*; *Op*, *Ophiomorpha*; *Pal*, *Paleodictyon*; *Ph*, *Phycosiphon*; *Pl*, *Planolites*; *Pr*, *Protopaleodictyon*; *Ro*, *Rosselia*; *Sc*, *Scolicia*; *Sp*, *Spirophyton*; *Th*, *Thalassinoides*; *Zo*, *Zoophycos*.

A basinward differentiation of the archetypical *Cruziana* ichnofacies – *Zoophycos* ichnofacies – *Nereites* ichnofacies must be manifest in the associated ichnological and sedimentological features. Yet the characterization of particular ichnofacies is hardly discernible in their transitions. The distal expression of the *Cruziana* ichnofacies can be interdigitated with the *Zoophycos* ichnofacies, and differentiation between the *Zoophycos* ichnofacies and the *Nereites* ichnofacies can be difficult in the case of fine-grained, distal turbidites, when sedimentation is more or less constant and low.


## 2. MATERIALS AND METHODS

The full section of Site U1385 from the IODP Expedition 339 (Expedition 339 Scientists, 2013a; Hernández-Molina *et al.*, 2013) was studied. This Site is located on the West Iberian margin (37°34.285'N, 10°7.562'W, Fig.2), on a spur, near the Core MD95-2042 (the “Shackleton Site”), a location with special relevance when studying millennial-scale climate variability over the last glacial cycles (Shackleton *et al.*, 2000, 2004) or interpreting global environmental conditions based on correlations with worldwide records (Expedition 339 Scientists, 2013b; Hodell *et al.*, 2013, 2015). The record of this site mainly comprises four cores from around 150 m penetration depth each (U1385A, U1385B, U1385D, and U1385E; total recovery of core U1385C is around 10 m),

representing continuous deposition from ~1.45 Ma ago to the Holocene (Expedition 339 Scientists, 2013b; Hodell *et al.*, 2013, 2015). Sediments are mainly composed by bioturbated calcareous muds and calcareous clays; primary sedimentary structures were not observed (Expedition 339 Scientists, 2013a, b).

High-resolution digital image treatment (Dorador & Rodríguez-Tovar, 2014, 2015; Dorador *et al.*, 2014a, b; Rodríguez-Tovar *et al.*, 2015a, b; Takashimizu *et al.*, 2015) was applied to original images, scanned with 500 dpi image resolution, of cores from Site U1385 (Expedition 339 Scientists, 2013c). These images were treated following the modifications proposed by Dorador *et al.* (2014a) to improve the visibility of trace fossils and ichnotaxonomic determination. This modification entails three image adjustments: 1) *levels*, 2) *brightness* and 3) *vibrance* (Dorador *et al.*, 2014a). The *Levels* adjustment involves stretching of the histogram according to colour extremes, modifying three parameters (black, grey and white). Black and white are established to represent the extreme values of the histogram, while the grey value is modified to make the image darker or lighter. The *Brightness* adjustment of reflected light can be used to modify two parameters, brightness and contrast. High values in both parameters mean a greater amount of reflected light and greater contrast of the image. Finally, the *vibrance* adjustment corrects the yellowish tone that is produced during execution of the two previous adjustments, using as values vibrance and saturation. After several modifications, the most positive results for the





**Figure 2.** Location of Site U1385, IODP Expedition 339, and lithologic log for the site.


selected adjustments were obtained for *levels* from 25 to 32 black, 0.43 to 0.75 grey and 92 to 100 white; *brightness* from -32 to -12 and contrast from 43 to 65; and *vibrance* from -7 to -11 for the vibrance parameter and from -12 to -18 for saturation values.

Occasionally, when ichnotaxonomical determination is particularly complicated due to similar geometries in different ichnotaxa, the high-resolution digital image method was supplemented by the quantification and comparison of pixel values in the infilling material of these ichnotaxa (Dorador & Rodríguez-Tovar, 2014).

### 3. RESULTS

#### 3.1. Ichnotaxa and ichnofacies characterization


The proposed methodology allowed for differentiation of eight ichnotaxa in the studied cores from Site U1385. Trace fossil assemblages, similar to the one differentiated by Rodríguez-Tovar & Dorador (2014), consist of abundant *Planolites*, common *Palaeophycus* and *Thalassinoides*, together with less abundant *Taenidium* and *Thalassinoides*-like structures, localized *Zoophycos* and *Chondrites*, and rare *Phycosiphon* and horizontal trails (?*Nereites*) (Fig. 3). *Chondrites* is observed as clusters of very small circular to elliptical spots and short tubes. *Palaeophycus* consists of circular to subcircular, unbranched lined cylindrical structures infilled by material having the same lithology as the host rock. *Phycosiphon* was registered as randomly oriented patches of horizontal, curved small lobes,



**Figure 3.** Schematic drawings, in 2D and 3D, of trace fossils identified in the study cores (following Wetzel & Uchman, 2012).

showing dark, fine-grained cylindrical to circular cores. *Planolites* appears as circular to subcircular unbranched cylindrical unlined tubular forms, filled with sediment of different material than that of the host sediment. *Taenidium* was found as horizontal to oblique, simple, straight to sinuous, unlined tubular meniscate structures. *Thalassinoides* are larger circular to subcircular oval spots and straight or slightly winding horizontal to oblique cylinders. *Thalassinoides*-like structures are similar to *Thalassinoides*, but show a variably developed irregular wall and diffuse shape. *Zoophycos* consists of repeated, horizontal to subhorizontal, spreite structures. Densely packed large black circular forms, enveloped by light sediments probably associated with horizontal trails, are tentatively assigned to ?*Nereites*.

Trace fossil assemblages, together with sedimentological features of the studied deposits, allowed us to identify the *Zoophycos* ichnofacies, especially well-represented in those intervals where *Zoophycos* is dominant. *Zoophycos*, locally accompanied by *Chondrites*, was seen to locally crosscut *Planolites* and *Thalassinoides*. Even though the *Zoophycos* ichnofacies could be clearly differentiated, observed trace fossil assemblages in some intervals of the studied cores could be related to the distal expression of the *Cruziana* ichnofacies and to the *Nereites* ichnofacies (Fig. 4). Intervals characterized by abundant *Planolites*, *Thalassinoides* and *Thalassinoides*-like structures, the subordinate presence of *Palaeophycus* and *Taenidium*, and rare *Chondrites* and *Zoophycos*, can be assigned to the distal expression of *Cruziana* ichnofacies, while, the local record of densely-packed, large, black circular forms enveloped by light sediments (?*Nereites*), may reveal the presence of grazing trails typical of the *Nereites* ichnofacies.



**Figure 4.** Examples of the three expressions of the *Zoophycos* ichnofacies identified in this study (archetypical *Zoophycos* ichnofacies, *Zoophycos* ichnofacies with elements of the distal *Cruziana* ichnofacies, and *Zoophycos* ichnofacies with elements of the *Nereites* ichnofacies). ?*Ne*, ?*Nereites*; *Pa*, *Palaeophycus*; *Pl*, *Planolites*; *Th*, *Thalassinoides*; *Th-l*, *Thalassinoides*-like structures; *Zo*, *Zoophycos*. Scale bar 1 cm.

#### 4. DISCUSSION

Sedimentological and ichnological data from Site U1385 are compatible with deposition under low-energy conditions in a deep-sea hemipelagic setting. The sediments in the studied cores mainly consists of mudstone (Expedition 339 Scientists, 2013b), with a relatively constant grain size, deposited by suspension fall-out. Trace fossil assemblages generally support good bottom- and pore-water conditions and organic matter availability, with only exceptional variations in these parameters (Rodríguez-Tovar & Dorador, 2014). Precise ichnofacies characterization might therefore be very useful to refine palaeoenvironmental interpretations.

As indicated above, generalized archetypical *Zoophycos* ichnofacies could be differentiated, but particular ichnotaxa may correspond to the distal expression of *Cruziana* ichnofacies (i.e., frequent *Thalassinoides* and *Thalassinoides*-like) or to the *Nereites* ichnofacies (i.e., ?*Nereites*). Before making any conclusive ichnofacies assignments, it is necessary to consider that: 1) these particular ichnotaxa are also registered in the *Zoophycos*

ichnofacies, 2) other typical elements of both the distal expression of the *Cruziana* and the *Nereites* ichnofacies are absent, and 3) the *Zoophycos* ichnofacies shows intergradation landward with the distal *Cruziana* ichnofacies and basinward with the *Nereites* ichnofacies. On this basis the *Zoophycos* ichnofacies can be considered as the only one registered in the studied succession, but with variations, including its archetypical expression and those showing elements also registered in the distal expression of *Cruziana* ichnofacies or in the *Nereites* ichnofacies.

The proposed discrimination among three distinct expressions of the *Zoophycos* ichnofacies is relevant when interpreting the depositional environment of the studied succession. Environmental parameters influencing tracemakers —sedimentation rate, energy conditions, sea level, oxygenation, and food supply, among others— vary with distance from shore. Thus, the distinct ichnofacies expressions could help recognize changing depositional conditions. A clear example would be the core section seen in Figure 5, corresponding to core U1385A-15H-3-A from Site U1385 of IODP Expedition 339. Overall,



**Figure 5.** Differentiation between the *Zoophycos* ichnofacies with elements of the distal expression of *Cruziana* ichnofacies and the archetypal *Zoophycos* ichnofacies in core U1385A-15H-3-A. *Ch*, *Chondrites*; *Ph*, *Phycosiphon*; *Pl*, *Planolites*; *Th*, *Thalassinoides*; *Zo*, *Zoophycos*. Scale bar 1 cm.

it has a mottled background ichnofabric, on which some discrete trace fossils can be observed. The trace fossil assemblage consists of dominant *Zoophycos* and subordinate *Chondrites*, *Planolites* and *Thalassinoides*, indicating the *Zoophycos* ichnofacies. In this generalized context, however, stratigraphical variations are recognized. From the bottom to top of the core *Zoophycos* progressively decreases in abundance and eventually disappears (in the upper part), while the abundance of *Planolites* and *Thalassinoides* increase significantly. This could be interpreted as a continuous change from the archetypal *Zoophycos* ichnofacies to the *Zoophycos* ichnofacies with elements of the distal expression of the *Cruziana* ichnofacies. This subtle change might be related to minor changes in the palaeoenvironmental conditions (e.g., slight increase in oxygenation, food content and/or deposition rate associated with more proximal depositional settings).

## 5. CONCLUSIONS

A high-resolution image procedure was applied to cores from Site U1385 (IODP Expedition 339) to improve characterization of deep-sea ichnofacies. This method is mainly based on the modification of three image adjustments: *levels*, *brightness* and *vibrance*, together with pixel analysis.

Eight ichnotaxa were recognized (*Chondrites*, *Palaeophycus*, *Phycosiphon*, *Planolites*, *Taenidium*, *Thalassinoides*, *Thalassinoides*-like structures and *Zoophycos*) along with undefined horizontal trails (?*Nereites*). Variations in trace fossil diversity and abundance occur throughout the succession.

Sedimentological and ichnological data indicate the dominance (exclusiveness) of the *Zoophycos* ichnofacies, but reveal three different expressions: 1) archetypal *Zoophycos* ichnofacies, characterized by dominance of *Zoophycos*, with subordinate *Chondrites*, *Planolites*, *Thalassinoides* and *Thalassinoides*-like structures; 2) *Zoophycos* ichnofacies with elements of the distal expression of the *Cruziana* ichnofacies, consisting of common *Planolites*, *Thalassinoides* and *Thalassinoides*-like structures; and 3) *Zoophycos* ichnofacies with elements of the *Nereites* ichnofacies, characterized by the presence of horizontal, grouped circular structures interpreted as grazing trails (?*Nereites*).



Vertical variations in the three expressions of the *Zoophycos* ichnofacies point to subtle changes in the depositional conditions potentially related to temporal variations in distance from shore, sedimentation rate, oxygenation and/or food content.

## ACKNOWLEDGEMENTS

This paper benefited from comments and suggestions by Drs Mayoral (Univ. Huelva) and Savrda (Auburn Univ.). The study has been funded by Project CGL2012-33281 (Secretaría de Estado de I+D+I, Spain) and a Research Grant from the Spanish Society of Palaeontology obtained by JD. The research of JD is funded through a pre-doctoral grant at the University of Granada.

## REFERENCES

- Bottjer, D.J., Droser, M.L. & Jablonski, D. 1987. Bathymetric trends in the history of trace fossils. In: *New Concepts in the Use of Biogenic Sedimentary Structures for Paleoenvironmental Interpretation* (ed. Bottjer, D.J.). Society of Economic Paleontologists and Mineralogists, Los Angeles, CA, 57-65.
- Bromley, R.G. 1990. *Trace fossils. Biology and Taphonomy*. Unwin Hyman, London.
- Bromley, R.G. 1996. *Trace fossils. Biology, Taphonomy and Applications*. Chapman & Hall, London.
- Buatois, L.A. & Mángano, M.G. 1995. The paleoenvironmental and paleoecological significance of the lacustrine *Mermia* ichnofacies: An archetypical subaqueous nonmarine trace fossil assemblage. *Ichnos*, 4 (2), 151-161.
- Buatois, L.A. & Mángano, M.G. 2011. *Ichnology: Organism-Substrate Interactions in Space and Time*. Cambridge University Press, Cambridge.
- Dorador, J. & Rodríguez-Tovar, F.J. 2014. A novel application of quantitative pixels analysis to trace fossil research in marine cores. *Palaaios*, 29, 533-538.
- Dorador, J. & Rodríguez-Tovar, F.J. 2015. Stratigraphic variation in ichnofabrics at the "Shackleton Site" (IODP Site U1385) on the Iberian Margin: Paleoenvironmental implications. *Marine Geology*, doi: 10.1016/j.margeo.2015.09.008.
- Dorador, J., Rodríguez-Tovar, F.J. & IODP Expedition 339 Scientists. 2014a. Digital image treatment applied to ichnological analysis of marine core sediments. *Facies*, 60 (1), 39-44.
- Dorador, J., Rodríguez-Tovar, F.J. & IODP Expedition 339 Scientists. 2014b. Quantitative estimation of bioturbation based on digital image analysis. *Marine Geology*, 349, 55-60.
- Expedition 339 Scientists. 2013a. Expedition 339 summary. In: *Proceedings of the Integrated Ocean Drilling Program, Tokyo* (eds. Stow, D.A.V., Hernández-Molina, F.J., Alvarez-Zarikian, C.A. & Expedition 339 Scientists). Integrated Ocean Drilling Program Management International, Inc.
- Expedition 339 Scientists. 2013b. Site U1385. In: *Proceedings of the Integrated Ocean Drilling Program, Tokyo* (eds. Stow, D.A.V., Hernández-Molina, F.J., Alvarez-Zarikian, C.A. & Expedition 339 Scientists). Integrated Ocean Drilling Program Management International, Inc.
- Expedition 339 Scientists. 2013c. Methods. In: *Proceedings of the Integrated Ocean Drilling Program, Tokyo* (eds. Stow, D.A.V., Hernández-Molina, F.J., Alvarez-Zarikian, C.A. & Expedition 339 Scientists). Integrated Ocean Drilling Program Management International, Inc.
- Frey, R.W. & Pemberton, S.G. 1985. Biogenic structures in outcrops and cores. I. Approaches to ichnology. *Bulletin of Canadian Petroleum Geology*, 33, 72-115.
- Heard, T.G. & Pickering, K.T. 2008. Trace fossils as diagnostic indicators of deep marine environments, Middle Eocene Ainsa Jaca basin, Spanish Pyrenees. *Sedimentology*, 55, 809-844.
- Hernández-Molina, F.J., Stow, D., Álvarez-Zarikian, C.A. & Expedition IODP 339 Scientists. 2013. IODP Expedition 339 in the Gulf of Cadiz and off West Iberia: decoding the environmental significance of the Mediterranean outflow water and its global influence. *Scientific Drilling*, 16, 1-11.
- Hodell, D.A., Lourens, L., Stow, D.A.V., Hernández-Molina, F.J., Alvarez-Zarikian, C. & Members, S.S.P. 2013. The "Shackleton Site" (IODP Site U1385) on the Iberian Margin. *Scientific Drilling*, 16, 13-19.
- Hodell, D.A., Lourens, L., Crowhurst, S., Konijnendijk, T., Tjallingii, R., Jiménez-Espejo, F., Skinner, L., Tzedakis, P.C. & Shackleton Site Project Members. 2015. A reference time scales for Site U1385 (Shackleton Site) on the SW Iberian Margin. *Global and Planetary Change*, 133, 49-64.
- Howard, J.D. 1968. X-ray radiography for examination of burrowing in sediments by marine invertebrate organisms. *Sedimentology*, 11, 249-258.
- Hubbard, S.M., MacEachern, J.A. & Bann, K.L. 2012. Slopes. In: *Trace Fossils as Indicators of Sedimentary Environments* (eds. Knaust, D. & Bromley, R.G.). Developments in Sedimentology 64, Elsevier, Amsterdam, 607-642.
- Joschko, M., Graff, O., Müller, P.C., Kotzke, K., Lindner, O., Pretschner, D.P. & Larink, O. 1991. A non-destructive method for the morphological assessment of earthworm burrow systems in three dimensions by X-ray computed tomography. *Biology and Fertility of Soils*, 11, 88-92.
- Knaust, D. 2009. Characterisation of a Campanian deep-sea fan system in the Norwegian Sea by means of ichnofabrics. *Marine and Petroleum Geology*, 26, 1119-1211.
- Löwemark, L. & Werner, F. 2001. Dating errors in high resolution stratigraphy: a detailed X-ray radiograph and AMS-14C study of *Zoophycos* Burrows. *Marine Geology*, 17, 191-198.
- MacEachern, J.A., Zaitlin, B.A. & Pemberton, S.G. 1999. A sharp-based sandstone of the Vinking Formation, Joffre Field, Alberta, Canada: criteria for recognition of



- transgressively incised shoreface complexes. *Journal of Sedimentary Research*, 69, 876-892.
- MacEachern, J.A., Bann, K.L., Pemberton, S.G. & Gingras, M.K. 2007. The Ichnofacies paradigm: High-resolution paleoenvironmental interpretation of the rock record. In: *Applied Ichnology* (eds. MacEachern, J.A., Bann, K.L., Gingras, M.K. & Pemberton, S.G.). Society for Sedimentary Geology Short Course Notes, 27-64.
- MacEachern, J.A., Bann, K.L., Gingras, M.K., Zonneveld, J.-P., Dashtgard, S.E. & Pemberton, S.G. 2012. The ichnofacies paradigm. In: *Trace fossils as indicators of sedimentary environments* (eds. Knaust, D. & Bromley, R.G.). *Developments in Sedimentology* 64, Elsevier, Amsterdam, 103-138.
- MacEachern, J.A., Pemberton, S.G., Gingras, M.K. & Bann, K.L. 2010. Ichnology and facies models. In: *Facies Models 4* (eds. James, N.O. & Dalrymple, R.W.). Geological Association of Canada, 19-58.
- Magwood, J.P.A. & Ekdale, A.A. 1994. Computer-aided analysis of visually complex ichnofabrics in deep-sea sediments. *Palaios*, 9, 102-115.
- Pemberton, S.G., MacEachern, J.A., Dashtgard, S.E., Bann, K.L., Gingras, M.K. & Zonneveld, J.-P. 2012. Shorefaces. In: *Trace fossils as indicators of sedimentary environments* (eds. Knaust, D. & Bromley, R.G.). *Developments in Sedimentology* 64, Elsevier, Amsterdam, 563-603.
- Rodríguez-Tovar, F.J. & Dorador, J. 2014. Ichnological analysis of Pleistocene sediments from the IODP Site U1385 "Shackleton Site" on the Iberian Margin: Approaching palaeoenvironmental conditions. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 409, 24-32.
- Rodríguez-Tovar, F.J. & Dorador, J. 2015. Ichnofabric characterization in cores: a method of digital image treatment. *Annales Societatis Geologorum Poloniae*, 85, 465-471.
- Rodríguez-Tovar, F.J., Dorador, J., Grunert, P. & Hodell, D. 2015b. Deep-sea trace fossil and benthic foraminiferal assemblages across glacial Terminations 1, 2 and 4 at the "Shackleton Site" (IODP Expedition 339, Site U1385). *Global and Planetary Change*, 133, 359-370.
- Rodríguez-Tovar, F.J., Dorador, J., Martín-García, G.M., Sierro, F.J., Flores, J.A. & Hodell, D.A. 2015a. Response of macrobenthic and foraminifer communities to changes in deep-sea environmental conditions from Marine Isotope Stage (MIS) 12 to 11 at the "Shackleton Site". *Global and Planetary Change*, 133, 176-187.
- Rosenberg, R., Davey, E., Gunnarsson, J., Norling, K. & Frank, M. 2007. Application of computer-aided tomography to visualize and quantify biogenic structures in marine sediments. *Marine Ecology Progress Series*, 331, 23-34.
- Seilacher, A. 1964. Biogenic sedimentary structures. In: *Approaches to Palaeoecology* (eds. Imbrie, J. & Newell, N.). John Wiley & Sons, Chichester, UK, 296-316.
- Seilacher, A. 1967. Bathymetry of trace fossils. *Marine Geology*, 5, 413-428.
- Seilacher, A. 1974. Flysch trace fossils: evolution of behavioural diversity in the deep-sea. *Neues Jahrbuch für Geologie und Paläontologie*, Monatshefte 1974, 223-245.
- Shackleton, N.J., Hall, M.A. & Vincent, E. 2000. Phase relationships between millennial-scale events 64,000–24,000 years ago. *Paleoceanography*, 15 (6), 565-569.
- Shackleton, N., Fairbanks, R., Chiu, T.C. & Parrenin, F. 2004. Absolute calibration of the Greenland time scale: implications for Antarctic time scales and for  $\delta^{14}\text{C}$ . *Quaternary Science Reviews*, 23 (14), 1513-1522.
- Takashimizu, Y., Kawamura, R., Rodríguez-Tovar, F.I., Dorador, J., Ducassou, E., Hernández-Molina, F.I., Stow, D.V.A. & Alvarez-Zarikian, C.A. 2015. Reworked Tsunami deposits by bottom currents: Circumstantial evidences from Late Pleistocene to Early Holocene in de Gulf of Cádiz. *Marine Geology*, doi:10.1016/j.margeo.2015.09.009.
- Uchman, A. 2009. The *Ophiomorpha rudis* ichnosubfacies of the *Nereites* ichnofacies: Characteristics and constraints. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 276 (1-4), 107-119.
- Uchman, A. & Wetzel, A. 2011. Deep-sea ichnology: The relationships between depositional environment and endobenthic organisms. In: *Deep-sea Sediments* (eds. Hüneke, H. & Mulder, T.). *Developments in Sedimentology* 63, Elsevier, Amsterdam, 517-556.
- Uchman, A. & Wetzel, A. 2012. Deep-Sea Fans. In: *Trace fossils as indicators of sedimentary environments* (eds. Knaust, D. & Bromley, R.G.). *Developments in Sedimentology* 64, Elsevier, Amsterdam, 643-671.
- Wetzel, A. 1991. Ecologic interpretation of deep-sea trace fossil communities. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 85 (1), 47-69.
- Wetzel, A. 2010. Deep-sea ichnology: Observations in modern sediments to interpret fossil counterparts. *Acta Geologica Polonica*, 60 (1), 125-138.
- Wetzel, A. & Uchman, A. 2012. Hemipelagic and Pelagic Basin Plains. In: *Trace fossils as indicators of sedimentary environments* (eds. Knaust, D. & Bromley, R.G.). *Developments in Sedimentology* 64, Elsevier, Amsterdam, 673-701.

